Please check the examination details below before entering your candidate information

Candidate surname	Other names
Solved by Mr. Pabitra	

Centre Number | Candidate Number

Pearson Edexcel International Advanced Level

Thursday 22 May 2025

Morning (Time: 1 hour 30 minutes)

Paper reference **WFM01/01**

Mathematics

International Advanced Subsidiary/Advanced Level
Further Pure Mathematics F1

You must have:
Mathematical Formulae and Statistical Tables (Yellow), calculator

Total Marks

Candidates may use any calculator allowed by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions:

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
  – *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information:

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 9 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
  – *use this as a guide as to how much time to spend on each question.*

Advice:

Solved by Mr. Pabitra | Math Tonic

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ▶

P76409A

©2025 Pearson Education Ltd.
Y:1/1/1/

P 7 6 4 0 9 A 0 1 3 6

Pearson

1: $$\mathbf{M} = \begin{pmatrix} 1 & a \\ 3 & -5 \end{pmatrix}$$

where a is a constant, $a \neq -\frac{5}{3}$

(a) Determine $\mathbf{M}^{-1}$ in terms of a.

(3)

Given that $\mathbf{M}^{-1} = 2\mathbf{M} + 8\mathbf{I}$ where $\mathbf{I}$ is the 2×2 identity matrix,

(b) determine the value of a.

(3)

(a)

$\det M = 1 \times (-5) - a \times 3 = -5 - 3a$

$$M^{-1} = \frac{1}{\det M}\begin{pmatrix} -5 & -a \\ -3 & 1 \end{pmatrix} = \frac{1}{-5-3a}\begin{pmatrix} -5 & -a \\ -3 & 1 \end{pmatrix}$$

$$\boxed{M^{-1} = \frac{1}{-5-3a}\begin{pmatrix} -5 & -a \\ -3 & 1 \end{pmatrix}}$$

(b)

$$2M + 8I = \begin{pmatrix} 2 & 2a \\ 6 & -10 \end{pmatrix} + \begin{pmatrix} 8 & 0 \\ 0 & 8 \end{pmatrix} = \begin{pmatrix} 10 & 2a \\ 6 & -2 \end{pmatrix}$$

Compare the top-left elements:

$$\frac{-5}{-5-3a} = 10$$

$$-5 = 10(-5-3a) = -50 - 30a$$

$$30a = -45$$

$$a = \frac{-45}{30} = -\frac{3}{2}$$

Check: $\det M = -5 + 4.5 = -0.5$, $\frac{-3}{-0.5} = 6$ and $\frac{1}{-0.5} = -2$ correct

$$\boxed{a = -\frac{3}{2}}$$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Solved by Mr. Pabitra

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 1 continued

Solved by Mr. Pabitra

Question 1 continued

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Solved by Mr. Pabitra

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 1 continued

(Total for Question 1 is 6 marks)

Solved by Mr. Pabitra

2:

$$\mathbf{P} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

The matrix **P** represents the transformation P.

(a) Describe P fully as a single geometrical transformation.

(2)

$$\mathbf{Q} = \begin{pmatrix} 5 & 0 \\ 0 & 5 \end{pmatrix}$$

The matrix **Q** represents the transformation Q.

(b) Describe Q fully as a single geometrical transformation.

(2)

The transformation R is a reflection in the line $y = -x$

(c) Write down the matrix representing R.

(1)

The transformation R maps the point $(4, 3)$ to the point A.

(d) Determine the coordinates of A.

(1)

The transformation P followed by the transformation Q maps the point B to the point A.

(e) Determine the coordinates of B.

(2)

(a) (1, 0) goes to (0, −1) and (0, 1) goes to (1, 0)
Rotation of 90° clockwise about the origin (0)

(b) Enlargement, scale factor 5, centre the origin (0)

(c) Reflection in $y = -x$: (1, 0) goes to (0, −1), (0, 1) goes to (−1, 0)

$R = \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}$

(d)

$\begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}\begin{pmatrix} 4 \\ 3 \end{pmatrix} = \begin{pmatrix} -3 \\ -4 \end{pmatrix}$ $A = (-3, -4)$

(e) P followed by Q is the matrix QP

$QP = \begin{pmatrix} 5 & 0 \\ 0 & 5 \end{pmatrix}\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 5 \\ -5 & 0 \end{pmatrix}$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Solved by Mr. Pabitra

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 2 continued

$$\begin{pmatrix} 0 & 5 \\ -5 & 0 \end{pmatrix}\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -3 \\ -4 \end{pmatrix}$$

$5y = -3$ so $y = -\frac{3}{5}$ $\quad -5x = -4$ so $x = \frac{4}{5}$

$$B = \left(\frac{4}{5}, -\frac{3}{5}\right)$$

MATH TONIC

Question 2 continued

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

MATH TONIC

Solved by Mr. Pabitra

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 2 continued

(Total for Question 2 is 8 marks)

Solved by Mr. Pabitra

3: **In this question you must show all stages of your working.**
Solutions relying entirely on calculator technology are not acceptable.

(i) $f(x) = x^2 + 5 - 8^{5x}$

Given that the equation $f(x) = 0$ has a single root, α, in the interval $[0, 1]$

use interval bisection to determine an interval of width 0.25 that contains α.

(3)

(ii) $g(x) = 3^{\sin x} - 3\cos x$

where x is in radians.

(a) Show that the equation $g(x) = 0$ has a root, β, in the interval $[4, 5]$

(2)

(b) Use linear interpolation once on the interval $[4, 5]$ to determine an approximation for β.
Give your answer to 4 significant figures.

(2)

(i) $f(x) = x^2 + 5 - 8^{5x}$

$f(0) = 0 + 5 - 1 = 4 > 0$ $\quad f(1) = 1 + 5 - 32\,768 = -32\,762 < 0$

$f(0.5) = 0.25 + 5 - 8^{2.5} = -175.769... < 0$ so α is in $[0, 0.5]$

$f(0.25) = 0.0625 + 5 - 8^{1.25} = -8.391... < 0$ so α is in $[0, 0.25]$

$\boxed{0 \leqslant \alpha \leqslant 0.25}$

(ii)(a) $g(x) = 3^{\sin x} - 3\cos x$

$g(4) = 3^{\sin 4} - 3\cos 4 = 2.396... > 0$

$g(5) = 3^{\sin 5} - 3\cos 5 = -0.502... < 0$

There is a change of sign and $g(x)$ is continuous on $[4, 5]$,
so $g(x) = 0$ has a root β in the interval $[4, 5]$.

(ii)(b) Linear interpolation:

$$\frac{\beta - 4}{2.396...} = \frac{5 - \beta}{0.502...}$$

$0.502...(\beta - 4) = 2.396...(5 - \beta)$

$\beta(0.502... + 2.396...) = 4 \times 0.502... + 5 \times 2.396...$

$$\beta = \frac{2.009... + 11.981...}{2.898...} = 4.8267...$$

$\boxed{\beta = 4.827 \text{ (4 s.f.)}}$

DO NOT WRITE IN THIS AREA DO NOT WRITE IN THIS AREA DO NOT WRITE IN THIS AREA

Solved by Mr. Pabitra

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 3 continued

Solved by Mr. Pabitra

Question 3 continued

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

MATH TONIC

Solved by Mr. Pabitra

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 3 continued

(Total for Question 3 is 7 marks)

Solved by Mr. Pabitra

4: The rectangular hyperbola C has equation $xy = 81$

The point $P\left(9t, \frac{9}{t}\right)$, $t \neq 0$, lies on C.

(a) Use calculus to show that the normal to C at P has equation

$$ty = t^3x + 9\left(1 - t^4\right)$$

(4)

The normal to C at P meets the y-axis at the point A.

(b) Determine the exact coordinates of A in terms of t.

(2)

Given that $t = \frac{1}{3}$

(c) find the exact area of triangle OPA, where O is the origin.

(2)

(a) $y = \frac{81}{x} = 81x^{-1}$ $\quad \frac{dy}{dx} = -81x^{-2} = -\frac{81}{x^2}$

At P, $x = 9t$: $\quad \frac{dy}{dx} = -\frac{81}{81t^2} = -\frac{1}{t^2}$ (gradient of tangent)

Gradient of normal $= -1 \div \left(-\frac{1}{t^2}\right) = t^2$

Normal: $y - \frac{9}{t} = t^2(x - 9t)$

Multiply by t: $\quad ty - 9 = t^3x - 9t^4$

$ty = t^3x + 9 - 9t^4$

$ty = t^3x + 9(1 - t^4)$ $\quad$ as required

(b) At the y-axis, $x = 0$: $\quad ty = 9(1 - t^4)$

$y = \frac{9(1 - t^4)}{t}$

$$A = \left(0, \frac{9(1 - t^4)}{t}\right)$$

(c) $t = \frac{1}{3}$: $\quad P = \left(9 \times \frac{1}{3}, 9 \div \frac{1}{3}\right) = (3, 27)$

y-coordinate of A $= 9\left(1 - \frac{1}{81}\right) \div \frac{1}{3} = 27 \times \frac{80}{81} = \frac{80}{3}$

Triangle OPA: base OA on the y-axis, height = x-coordinate of P = 3

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Solved by Mr. Pabitra

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 4 continued

Area $= \frac{1}{2} \times \frac{80}{3} \times 3 = 40$

Area = 40

MATH TONIC

Solved by Mr. Pabitra

Question 4 continued

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

MATH TONIC

Solved by Mr. Pabitra

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 4 continued

(Total for Question 4 is 8 marks)

Solved by Mr. Pabitra

5: The complex number z_1 is given by

$$z_1 = r\left(\cos\frac{7\pi}{6} + i\sin\frac{7\pi}{6}\right)$$

where r is a positive constant.

The complex number z_2 has modulus 5

Given that $|z_1 z_2| = 15$

(a) state the value of r.

(1)

Given further that $z_1 + z_2$ is a real number,

(b) determine the possible complex numbers z_2 in the form $a + bi$ where a and b are constants.

(5)

(c) Show z_1 and the possible complex numbers z_2 on a single Argand diagram.

(2)

(a) $|z_1 z_2| = |z_1| \times |z_2| = r \times 5 = 15$

$r = 3$

(b) $z_1 = 3\left(\cos\frac{7\pi}{6} + i\sin\frac{7\pi}{6}\right) = -\frac{3\sqrt{3}}{2} - \frac{3}{2}i$

Let $z_2 = a + bi$. $z_1 + z_2$ is real, so the imaginary parts cancel:

$b - \frac{3}{2} = 0, \quad b = \frac{3}{2}$

$|z_2| = 5: \quad a^2 + \left(\frac{3}{2}\right)^2 = 25$

$a^2 = 25 - \frac{9}{4} = \frac{91}{4}$

$a = \pm\frac{\sqrt{91}}{2}$

$z_2 = \frac{\sqrt{91}}{2} + \frac{3}{2}i \quad \text{or} \quad z_2 = -\frac{\sqrt{91}}{2} + \frac{3}{2}i$


DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Solved by Mr. Pabitra

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 5 continued

(c) Argand diagram



MATH TONIC

Solved by Mr. Pabitra

Question 5 continued

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

MATH TONIC

Solved by Mr. Pabitra

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 5 continued

(Total for Question 5 is 8 marks)

Solved by Mr. Pabitra

6: $f(x) = 3x^2 + kx - 5$

where k is a constant.

The equation $f(x) = 0$ has roots α and β.

(a) State the value of $\alpha\beta$.

(1)

Given that $\alpha + \beta = 9\alpha\beta$

(b) determine the value of k.

(2)

(c) By first expanding $(\alpha + \beta)^3$ prove that

$$\alpha^3 + \beta^3 = (\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta)$$

(2)

Without solving the equation $f(x) = 0$

(d) find a quadratic equation with integer coefficients that has roots

$(\alpha^2 + \beta)$ and $(\alpha + \beta^2)$

(6)

(a) $\alpha\beta = \frac{c}{a} = \frac{-5}{3}$

$\boxed{\alpha\beta = -\frac{5}{3}}$

(b) $\alpha + \beta = -\frac{k}{3}$ $\quad -\frac{k}{3} = 9 \times \left(-\frac{5}{3}\right) = -15$

$\boxed{k = 45}$

(c) $(\alpha + \beta)^3 = \alpha^3 + 3\alpha^2\beta + 3\alpha\beta^2 + \beta^3$

$= \alpha^3 + \beta^3 + 3\alpha\beta(\alpha + \beta)$

so $\alpha^3 + \beta^3 = (\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta)$ as required

(d) $\alpha + \beta = -\frac{45}{3} = -15$, $\quad \alpha\beta = -\frac{5}{3}$

Sum of new roots: $(\alpha^2 + \beta) + (\alpha + \beta^2) = (\alpha^2 + \beta^2) + (\alpha + \beta)$

$= (\alpha + \beta)^2 - 2\alpha\beta + (\alpha + \beta) = 225 + \frac{10}{3} - 15 = \frac{640}{3}$

Product: $(\alpha^2 + \beta)(\alpha + \beta^2) = \alpha^3 + \alpha^2\beta^2 + \alpha\beta + \beta^3$

$= (\alpha^3 + \beta^3) + (\alpha\beta)^2 + \alpha\beta$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Solved by Mr. Pabitra

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 6 continued

$\alpha^3 + \beta^3 = (-15)^3 - 3(-\frac{5}{3})(-15) = -3375 - 75 = -3450$

Product $= -3450 + \frac{25}{9} - \frac{5}{3} = -\frac{31040}{9}$

Equation: $x^2 - \frac{640}{3}x - \frac{31040}{9} = 0$ $\quad (\times 9)$

$$\boxed{9x^2 - 1920x - 31\,040 = 0}$$

MATH TONIC

Question 6 continued

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

MATH TONIC

Solved by Mr. Pabitra

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 6 continued

(Total for Question 6 is 11 marks)

Solved by Mr. Pabitra

7: $f(z) = Pz^4 - 36z^3 + Qz^2 + 192z + 68$

where P and Q are real constants.

Given that $3 + 5i$ is a root of the equation $f(z) = 0$

(a) write down another root of the equation $f(z) = 0$

(1)

(b) Hence determine a quadratic factor, with real coefficients, of $f(z)$

(2)

(c) Determine the value of P and the value of Q.

(5)

Without using a calculator,

(d) determine the other roots of the equation $f(z) = 0$ giving your answers in simplest form.

(2)

(a) Real coefficients, so the complex conjugate is also a root:

$\boxed{3 - 5i}$

(b) Sum $= (3 + 5i) + (3 - 5i) = 6$, Product $= (3 + 5i)(3 - 5i) = 9 + 25 = 34$

$\boxed{z^2 - 6z + 34}$

(c) $f(z) = (z^2 - 6z + 34)(az^2 + bz + c)$

Constant term: $34c = 68$, $c = 2$

z term: $34b - 6c = 192$ $34b - 12 = 192$, $b = 6$

z^3 term: $b - 6a = -36$ $6 - 6a = -36$, $a = 7$

z^4 term: $P = a = 7$

z^2 term: $Q = c - 6b + 34a = 2 - 36 + 238 = 204$

$\boxed{P = 7, \quad Q = 204}$

(d) Other factor: $7z^2 + 6z + 2 = 0$

$$z = \frac{-6 \pm \sqrt{36 - 4 \times 7 \times 2}}{14} = \frac{-6 \pm \sqrt{-20}}{14} = \frac{-6 \pm 2\sqrt{5}i}{14}$$

$$\boxed{z = -\frac{3}{7} \pm \frac{\sqrt{5}}{7}i}$$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Solved by Mr. Pabitra

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 7 continued

Solved by Mr. Pabitra

Question 7 continued

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

MATH TONIC

Solved by Mr. Pabitra

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 7 continued

(Total for Question 7 is 10 marks)

Solved by Mr. Pabitra

8: (a) Using the standard summation formulae show that

$$\sum_{r=1}^{2n}\left(2r^2-1\right)=\frac{4}{3}n(n+1)(an+b)$$

where a and b are integers to be determined.

(5)

(b) Prove by induction that, for $n \in \mathbb{N}$

$$\sum_{r=1}^{n} r(3r-2)^2=\frac{n^2(n+1)(9n-7)}{4}$$

(5)

Using the results from parts (a) and (b) and showing all stages of your working,

(c) determine the value of n for which

$$8\sum_{r=1}^{n} r(3r-2)^2=15\sum_{r=1}^{2n}\left(2r^2-1\right)$$

(*Solutions relying entirely on calculator technology are not acceptable.*)

(3)

(a) $\sum_{r=1}^{2n}(2r^2-1)=2\sum_{r=1}^{2n}r^2-\sum_{r=1}^{2n}1$

$=2\times\frac{2n}{6}(2n+1)(4n+1)-2n$

$=\frac{2n}{3}[(2n+1)(4n+1)-3]=\frac{2n}{3}(8n^2+6n-2)$

$=\frac{4n}{3}(4n^2+3n-1)=\frac{4}{3}n(n+1)(4n-1)$

$a=4, \quad b=-1$

(b) $n=1$: LHS $=1\times(3-2)^2=1$ RHS $=\frac{1^2\times2\times2}{4}=1$ so true for $n=1$

Assume true for $n=k$: $\sum_{r=1}^{k}r(3r-2)^2=\frac{k^2(k+1)(9k-7)}{4}$

Then $\sum_{r=1}^{k+1}r(3r-2)^2=\frac{k^2(k+1)(9k-7)}{4}+(k+1)(3k+1)^2$

$=\frac{k+1}{4}[9k^3-7k^2+4(9k^2+6k+1)]=\frac{k+1}{4}(9k^3+29k^2+24k+4)$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Solved by Mr. Pabitra

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 8 continued

$= \frac{k+1}{4}(k+1)(9k^2+20k+4) = \frac{(k+1)^2(k+2)(9k+2)}{4}$

$= \frac{(k+1)^2((k+1)+1)(9(k+1)-7)}{4}$ which is the formula with $n = k+1$

It is true for $n = 1$, and if it is true for $n = k$ then it is true for $n = k+1$.

So by mathematical induction it is true for all positive integers n.

(c) $8 \times \frac{n^2(n+1)(9n-7)}{4} = 15 \times \frac{4}{3} n(n+1)(4n-1)$

$2n^2(n+1)(9n-7) = 20n(n+1)(4n-1)$

Divide by $2n(n+1)$: $n(9n-7) = 10(4n-1)$

$9n^2 - 7n - 40n + 10 = 0$ so $9n^2 - 47n + 10 = 0$

$(9n-2)(n-5) = 0$ $n = \frac{2}{9}$ (not an integer) or $n = 5$

$\boxed{n = 5}$

Solved by Mr. Pabitra

Question 8 continued

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

MATH TONIC

Solved by Mr. Pabitra

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 8 continued

(Total for Question 8 is 13 marks)

Solved by Mr. Pabitra

9:




Figure 1

Figure 1 shows a sketch of a parabola C.

The point P lies on C at a distance 5 units from the focus of C.

The points Q and R lie on the directrix of C, where $PQ = PR = 10$ units.

Determine the exact length of QR giving the answer in simplest form.

(4)

For a parabola, every point is the same distance from the focus and the directrix.
P is 5 from the focus, so the perpendicular distance from P to the directrix is 5.
Let M be the foot of this perpendicular on the directrix: PM = 5
Triangle PMQ is right-angled at M: $QM = \sqrt{PQ^2 - PM^2} = \sqrt{10^2 - 5^2} = \sqrt{75} = 5\sqrt{3}$
By symmetry $RM = QM = 5\sqrt{3}$
$QR = 2 \times 5\sqrt{3} = 10\sqrt{3}$

$\boxed{QR = 10\sqrt{3} \text{ units}}$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Solved by Mr. Pabitra

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 9 continued

Solved by Mr. Pabitra

Question 9 continued

(Total for Question 9 is 4 marks)

TOTAL FOR PAPER IS 75 MARKS

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Solved by Mr. Pabitra